


# **ProFoldin**

# P. aeruginosa MurF Assay Kits

P. aeruginosa MurF Assay Kit Plus-100 P. aeruginosa MurF Assay Kit Plus-500 Catalog No. MURF100KP Catalog No. MURF500KP

#### INTRODUCTION

MurF is the enzyme that catalyzes the last step in synthesis of UDP-MurNAc-pentapeptide in the pathway of peptidoglycan biosynthesis in bacteria. It is an essential enzyme and attractive target for anti-bacterial drug discovery. MurF adds a dipeptide D-Ala-D-Ala onto the MurE product UDP-MurNAc-tripeptide. The ligation reaction is coupled to the hydrolysis of ATP forming ADP and inorganic phosphate.



**UDP-MurNAc-tripeptide** 

D-Ala-D-Ala

**UDP-MurNAc-pentapeptide** 

The **P. aeruginosa** MurF Assay is based on measurement of the inorganic phosphate generated from the MurF reaction. The inorganic phosphate is detected by light absorbance at 650 nm. The assay reactions and detection can be performed by using 384-well or 96-well assay plates. Alternatively, the assay reaction can be carried out in Eppendorf tubes and the signal is measured using a cuvette. The high throughput assay can be used for screening inhibitors of *E.coli* MurF in drug discovery research. It may also be used for characterization of *E.coli* MurF.

The *P. aeruginosa* MurF Assay Kit Plus-100 (Catalog No. MURF100KP) contains the reagents for 100 assays in a 384-well plate assay format including 400 µl of 10 x Buffer, 33 µl of 100 x UDP-MurNAc-tripeptide (UMAG-DAP), 33 µl of 100 x D-Ala-D-Ala, 33 µl of 100 x ATP, 33 µl of 100 x *P. aeruginosa* MurF (2000 nM) and 5 ml of Dye MPA3000 for phosphate detection.

The *P. aeruginosa* MurF Assay Kit Plus-500 (Catalog No. MURF500KP) contains the reagents for 500 assays in a 384-well plate assay format including 2000 μl of 10 x Buffer, 170 μl of 100 x UDP-MurNAc-tripeptide (UMAG-DAP), 170 μl of 100 x D-Ala-D-Ala, 170 μl of 100 x ATP, 170 μl of 100 x *P. aeruginosa* MurF (2000 nM) and 25 ml of Dye MPA3000 for phosphate detection.

#### ASSAY PROTOCOL

The following assay protocol is based on the 384-well plate assay format. The reaction volume is 30 µl

#### ProFoldin

10 Technology Drive, Suite 40, Number 188 Hudson, MA 01749-2791 USA Phone: (508) 735-2539 FAX: (508) 845-9258

info@profoldin.com

www.profoldin.com

# **INSTRUCTIONS**

and the final assay volume is 75 µl. For 96-well plate assays, the reaction volume is 60 µl and the final assay volume is 150 µl. For detection using a cuvette, the reaction volume is 400 µl and the final assay volume is 1000 µl.

#### 1. Reagent preparation:

For each 10 assay reactions,

- (1) Prepare 297 μl of premix composed of 261 μl of H<sub>2</sub>O, 33 μl of 10 x Buffer and 3.3 μl of 100 x P. aeruginosa MurF.
- (2) Prepare 33 µl of 10 x Enzyme substrate by mixing 3.3 µl of 100 x UMAG-DAP, 3.3 µl of 100 x D-Ala-D-Ala, 3.3 µl of 100 x ATP, and 23.1 µl of water.

### 2. Reaction:

Mix 27 µl of the premix with 3 µl of the 10 x Enzyme substrate in each well. Incubate the reaction mixture at 37°C for 60 min.

#### 3. Detection:

Add 45 µl of the Dye MPA3000 into the 30 µl of the reaction mixture. Incubate for 5 min. Measure the light absorbance at 650 nm.

## **Assay Protocol for enzyme inhibition**

The assay can be optimized in terms of assay window, assay linearity and sensitivity to competitive inhibitors. ProFoldin offers HTS assay development service. For more information, please visit our website at http://www.profoldin.com/services.html.

### **Related Products**

E. coli MurF Assay Kit Plus-100	MURF100KE
S. aureus MurF Assay Kit Plus-100	MURF100KS
S. pneumoniae MurF Assay Kit Plus-100	MURF100KN
E. coli MurA Assay Kit Plus-100	MURA100KE
E. coli MurC Assay Kit Plus-100	MURC100KE
S. pneumoniae MurC Assay Kit Plus-100	MURC100KN
E. coli MurD Assay Kit Plus-100	MURD100KE
P. aeruginosa MurD Assay Kit Plus-100	MURD100KP
S. aureus MurD Assay Kit Plus-100	MURD100KS
E. coli MurE Assay Kit Plus-100	MURE100KE
E. coli GlmU Assay Kit Plus-100	GLU100KE
E. coli D-alanine ligase Assay Kit Plus-100	DDA100KE

# More information of drug targets and enzyme assays

For more information of drug targets and enzyme assays, please visit www.profoldin.com or send emails to info@profoldin.com.