



ProFoldin

10 Technology Drive, Suite 40, Number 188

Hudson, MA 01749-2791 USA

Tel: (508) 735-2539 FAX: (508) 845-9258

www.profoldin.com

info@profoldin.com



INSTRUCTIONS

ProFoldin

P. aeruginosa DNA Gyrase

P. aeruginosa DNA Gyrase – for 100 assays Catalog No. TOP2-100PA

| | |
|------------------------|--|
| Protein construct: | Wild-type <i>P. aeruginosa</i> DNA gyrase, the (gyrA) ₂ (gyrB) ₂ complex, composed of the gyrA subunit (MW 103 kDa) and gyrB subunit (92 kDa) purified from a bacterial expression system. |
| MW: | 390 kDa |
| Enzyme concentration: | 2 μM |
| Enzyme activity assay: | The DNA gyrase DNA supercoiling activity is measured by using the DNA Topoisomerase II (Gyrase) Assay Kit (Catalog No. DSA020K) |
| Storage temperature: | -20 or -80°C. Do not freeze-and-thaw repeatedly. |
| Enzyme dilution: | Use the 1 x assay to dilute the enzyme just before the assay. Do not store diluted enzyme solution |



The *P. aeruginosa* DNA Topoisomerase II (Gyrase) Assay Kit Plus-100 (Catalog No. TOP2-100PA) includes 50 μl 100 x *P. aeruginosa* gyrase (2 μM). It is for 100 assays.

Assay Protocol

The following assay protocol is based on a 96-well assay plate format using a standard black 96-well plate (Greiner 655076). For 384-well plate assays using a standard black 384-well plate (Matrix 4318), please reduce the reagent volumes proportionally.



ProFoldin

10 Technology Drive, Suite 40, Number 188
Hudson, MA 01749-2791 USA
Tel: (508) 735-2539 FAX: (508) 845-9258
www.profoldin.com
info@profoldin.com

INSTRUCTIONS

1. Reaction:

The total volume of each reaction mixture is 40 μ l including: 24 μ l of H₂O, 4 μ l of 10 x buffer, 4 μ l of 10 x relaxed DNA (250 μ g/ml relaxed plasmid DNA), 4 μ l of 10 x enzyme, 4 μ l of 10 mM ATP. Incubate the reaction mixture at 37°C for 60 min.

Note: The final concentrations are 20 mM Tris-HCl, pH 8, 35 mM NH₄OAc, 4.6 % glycerol, 1 mM DTT, 0.005% Brij35, 8 mM MgCl₂, 25 μ g/ml relaxed plasmid DNA, 1 mM ATP and 20 nM topoisomerase II. The 10 x enzyme is prepared by dilution of the 100x enzyme in 1x assay buffer. A negative control reaction can be the reaction mixture without addition of ATP. Magnesium is essential for the reaction and assay. EDTA should be avoided.

2. Assay

- (1) Make 1 x H19 Dilution Buffer by dilution of the 10 x H19 Dilution Buffer 10-fold with water. Freshly prepare the H19 dye by dilution of 1 μ l the 1500 x H19 dye stock solution with 1.5 ml of 1 x H19 Dilution Buffer (1500 x dilution). Mix the solution evenly by inverting the tube a few times.
- (2) Mix 250 μ l of the freshly prepared H19 dye with each reaction solution (40 μ l). Incubate the mixture at room temperature for 5 min.
- (3) Measure the fluorescence intensity at 535 nm using the excitation wavelength at 485 nm.

Related Products:

| | |
|--|-----------------------|
| DNA Topoisomerase II (Gyrase) Assay Kit | Catalog No. DSA020K |
| <i>P. aeruginosa</i> DNA Gyrase Assay Kit Plus-100 | Catalog No. DSA100KP |
| H19 Dye for DNA Relaxation and Supercoiling Assays | Catalog No. DSA1000D |
| <i>E. coli</i> DNA Gyrase Assay Kit Plus-100 | Catalog No. DSA100KE |
| <i>S. aureus</i> DNA Gyrase Assay Kit Plus-100 | Catalog No. DSA100KSE |
| Human Topoisomerase II DNA Decatenation Assay Kit Plus-100 | Catalog No. HDC100KE |
| Human Topoisomerase I DNA Relaxation Assay Kit Plus-100 | Catalog No. HRA100KE |
| 96-Well <i>E. coli</i> Topoisomerase I DNA Decatenation Assay Plus | Catalog No. T1DD96KE |
| 96-Well <i>E. coli</i> Topo IV DNA Decatenation Assay Plus | Catalog No. EDD96KE |

Publications

1. Asha M.K. et al, In vitro anti-Helicobacter pylori activity of a flavonoid rich extract of Glycyrrhiza glabra and its probable mechanisms of action, Journal of Ethnopharmacology, Vol.145(2), pp.581-586 (2013).
2. Mora-Pale M et al, Antimicrobial mechanism of resveratrol-trans-dihydrodimer produced from peroxidase-catalyzed oxidation of resveratrol. Biotechnol Bioeng. Vol 112, pp2417–2428 (2015).

For more information of DNA topoisomerase assays and assays for more drug targets and enzymes, please visit www.profoldin.com or send emails to info@profoldin.com.